

SOME REMARKS ON THE DERIVED CATEGORIES OF COHERENT SHEAVES ON HOMOGENEOUS SPACES

ALEXANDER SAMOKHIN

ABSTRACT. In this paper we prove first a general theorem on semiorthogonal decompositions in derived categories of coherent sheaves for flat families over a smooth base. We then show that the derived categories of coherent sheaves on flag varieties of classical type are generated by complete exceptional collections. Finally, we find complete exceptional collections in the derived categories of some homogeneous spaces of the symplectic groups of small rank.

1. INTRODUCTION

The study of derived categories of coherent sheaves on algebraic varieties dates back to the late 70's when A. Beilinson described the derived categories of projective spaces in [1]. Since then this field has experienced a great development that put it in the forefront of modern algebraic geometry. At the beginning of the 80's it was discovered that derived categories of coherent sheaves appeared in a variety of areas of mathematics: e.g., in representation theory of associative algebras, and in the study of moduli spaces, to name a few. Other links were found later, of which the homological mirror conjecture ([10]), and reinterpretation of the minimal model program in birational geometry in terms of semiorthogonal decompositions in derived categories ([5]) are, perhaps, the most important. Another profound connection between representations of semisimple Lie algebras in positive characteristic and the derived categories of coherent sheaves on Springer fibers of the Grothendieck–Springer resolutions of nilpotent cones — in particular, on homogeneous spaces — was established recently in [2].

In fact, homogeneous spaces of semisimple Lie groups were the first varieties studied in the context of derived categories of coherent sheaves. Next to the paper [1] was a series of papers by M. Kapranov who developed substantially the approach from *loc.cit.*. His work culminated in the article [9], where the derived categories of coherent sheaves on homogeneous spaces of Lie groups of type $\mathbf{A}_n$ and on quadrics were described. Far back in the late 80's, these facts, together with other evidences, led experts to believe that the derived category of any homogeneous space has a description similar to that from [1] and [9], i.e. that such a category is generated by an exceptional collection. More recently, the results of [4] indicate that derived categories of coherent sheaves on varieties admitting pavings by affine spaces (of which homogeneous spaces are a particular class) should have complete exceptional collections. Such collections are a very particular case of semiorthogonal decompositions. Over more than fifteen years, which have passed since the appearance of [9], a great many powerful techniques were invented to study semiorthogonal decompositions in derived categories. Despite this, explicit complete exceptional collections on arbitrary homogeneous spaces of semisimple Lie groups remain an unsolved problem up-to-date.

We limit ourselves to indicating one source of our interest in these collections. As was shown by S. Mukai ([16]) in his classification of prime Fano threefolds with $b_2 = 1$, such threefolds are

This work was supported in part by the French Government Fellowship and the RFFI award No. 02-01-22005.

closely connected to homogeneous spaces. In particular, some of these threefolds can be obtained as linear sections of appropriate homogeneous spaces. It is important to construct semiorthogonal decompositions in the derived categories of prime Fano threefolds for a number of reasons; however, these categories are fairly complicated objects. A recent phenomenon, discovered and studied by A. Kuznetsov in a series of papers ([11], [12], [13]), is that semiorthogonal decompositions in the derived categories of linear sections of homogeneous spaces turned out to be intrinsically related to those in the derived categories of linear sections of projectively dual varieties. The approach developed in *loc.cit.* permits, in particular, to describe in a uniform way the derived categories of linear sections of some homogeneous spaces. To make it work one needs to know the structure of derived category of a homogeneous space one starts with.

In the present paper we take some further steps towards the problem of finding complete exceptional collections on homogeneous spaces. In Section 3 we prove Theorem 3.1 on semiorthogonal decompositions for flat families over a smooth scheme. The assumptions of Theorem 3.1, however, are restrictive enough and, as a consequence, this theorem does not cover such important case as flat families of quadrics. Happily, semiorthogonal decompositions for such families were thoroughly studied in a recent paper [13]. Relying mainly on the results of *loc.cit.*, in Theorem 4.1 we show that the derived categories of flag varieties $\mathbf{G}/\mathbf{B}$, where $\mathbf{G}$ is a semisimple Lie group of classical type, are generated by exceptional collections. Note that Theorem 4.1 (for classical groups other than groups of type $\mathbf{A}_n$, which were treated in the paper [9] cited above) must have been known to experts for a long time; however, there were no references to this fact. In the final part of the paper we consider homogeneous spaces of the symplectic groups of small rank, and prove in Theorem 5.1 that the derived category of homogeneous space $\mathbf{Sp}_6/\mathbf{P}$, where $\mathbf{P}$ corresponds to the middle node of the Dynkin diagram of type $\mathbf{C}_3$, has a complete exceptional collection. This gives, together with Theorem 3.1 and the main result of [19], a complete description of the derived categories of homogeneous spaces of the group $\mathbf{Sp}_6$ and of half a number of homogeneous spaces of the group $\mathbf{Sp}_8$. Note also that the homogeneous space $\mathbf{Sp}_6/\mathbf{P}$ in Theorem 5.1 is a smooth hyperplane section of the grassmannian $\mathrm{Gr}_{2,6}$. Recently, the derived categories of hyperplane sections of the grassmannians $\mathrm{Gr}_{2,2n}$ for arbitrary n were treated independently by A. Kuznetsov in [14].

Finally, let us remark that a totally different approach to the above problems, which is inspired by the work [2], is discussed in a forthcoming paper [20].

NOTATION

All schemes are assumed to be smooth over an algebraically closed field $\mathbf{k}$ of characteristic zero. For a proper scheme X let $D^b(X)$ denote the bounded derived category of coherent sheaves on X . For an object $\mathcal{E}$ of $D^b(X)$ we denote $\mathcal{E}[n]$ the object $\mathcal{E}$ shifted by n . Given a morphism $f: X \rightarrow Y$, we denote the derived functors of push-forward and pull-back functors by f_* and f^* , respectively. Similarly, the derived tensor product is denoted simply by $\otimes$. For two objects A and B of $D^b(X)$ let $\mathcal{H}om_X^\bullet(A, B)$ denote the object $\mathcal{R}Hom_X(A, B) \in D^b(X)$ and let $Hom_X^\bullet(A, B)$ be the complex of $\mathbf{k}$ -vector spaces $RHom_{D^b(X)}(A, B)$. Given a scheme S , points of S always mean closed points. For a scheme S , and a vector bundle $\mathcal{E}$ over S , the projectivization of $\mathcal{E}$ is denoted $\mathbb{P}_S(\mathcal{E})$.

2. PRELIMINARIES

We recall here some basic definitions. The material below is taken from [3].

Throughout this section we fix a triangulated $\mathbf{k}$ -linear category $\mathbf{D}$, equipped with a shift functor $[1]: \mathbf{D} \rightarrow \mathbf{D}$. For two objects $A, B \in \mathbf{D}$ let $\text{Hom}_{\mathbf{D}}^{\bullet}(A, B)$ be the graded $\mathbf{k}$ -vector space $\bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\mathbf{D}}(A, B[i])$. Let $\mathbf{A} \subset \mathbf{D}$ be a full triangulated subcategory, that is a full subcategory of $\mathbf{D}$, which is closed under shifts and taking cones.

Definition 2.1. The right orthogonal $\mathbf{A}^{\perp} \subset \mathbf{D}$ is defined to be the full subcategory

$$(2.1) \quad \mathbf{A}^{\perp} = \{B \in \mathbf{D} : \text{Hom}_{\mathbf{D}}(A, B) = 0\}$$

for all $A \in \mathbf{A}$. The left orthogonal ${}^{\perp}\mathbf{A}$ is defined similarly.

Definition 2.2. A full triangulated subcategory $\mathbf{A}$ of $\mathbf{D}$ is called *right admissible* if the inclusion functor $\mathbf{A} \hookrightarrow \mathbf{D}$ has a right adjoint. Similarly, $\mathbf{A}$ is called *left admissible* if the inclusion functor has a left adjoint. Finally, $\mathbf{A}$ is *admissible* if it is both right and left admissible.

It is shown ([3], Lemma 3.1) that if a full triangulated category $\mathbf{A} \subset \mathbf{D}$ is right admissible then every object $X \in \mathbf{D}$ fits into a distinguished triangle

$$(2.2) \quad \dots \longrightarrow Y \longrightarrow X \longrightarrow Z \longrightarrow Y[1] \longrightarrow \dots$$

with $Y \in \mathbf{A}$ and $Z \in \mathbf{A}^{\perp}$. One then says that there is a semiorthogonal decomposition of $\mathbf{D}$ into the subcategories $(\mathbf{A}^{\perp}, \mathbf{A})$. More generally, assume given a sequence of full triangulated subcategories $\mathbf{A}_1, \dots, \mathbf{A}_n \subset \mathbf{D}$. Denote $\langle \mathbf{A}_1, \dots, \mathbf{A}_n \rangle$ the triangulated subcategory of $\mathbf{D}$ generated by $\mathbf{A}_1, \dots, \mathbf{A}_n$.

Definition 2.3. A sequence $(\mathbf{A}_1, \dots, \mathbf{A}_n)$ of admissible subcategories of $\mathbf{D}$ is called *semiorthogonal* if $\mathbf{A}_i \subset \mathbf{A}_j^{\perp}$ for $1 \leq i < j \leq n$. The sequence $(\mathbf{A}_1, \dots, \mathbf{A}_n)$ is called a *semiorthogonal decomposition* of $\mathbf{D}$ if $\langle \mathbf{A}_1, \dots, \mathbf{A}_n \rangle^{\perp} = 0$, that is $\mathbf{D} = \langle \mathbf{A}_1, \dots, \mathbf{A}_n \rangle$.

Definition 2.4. An object $E \in \mathbf{D}$ is said to be exceptional if there is an isomorphism of graded $\mathbf{k}$ -algebras

$$(2.3) \quad \text{Hom}_{\mathbf{D}}^{\bullet}(E, E) = \mathbf{k}.$$

A collection of exceptional objects $(E_0, \dots, E_n)$ in $\mathbf{D}$ is called exceptional if for $1 \leq i < j \leq n$ one has

$$(2.4) \quad \text{Hom}_{\mathbf{D}}^{\bullet}(E_j, E_i) = 0.$$

Denote $\langle E_0, \dots, E_n \rangle \subset \mathbf{D}$ the full triangulated subcategory generated by the objects $E_0, \dots, E_n$. One proves ([3], Theorem 3.2) that such a category is admissible. The collection $(E_0, \dots, E_n)$ in $\mathbf{D}$ is said to be *complete* if $\langle E_0, \dots, E_n \rangle^{\perp} = 0$, in other words $\mathbf{D} = \langle E_0, \dots, E_n \rangle$.

Definition 2.5. Let (E_0, E_1) be an exceptional pair in $\mathbf{D}$, i.e. an exceptional collection of two elements. The left mutation of (E_0, E_1) is a pair $(L_{E_0}E_1, E_0)$, where the object $L_{E_0}E_1$ is defined to be a cone of the triangle

$$(2.5) \quad \dots \longrightarrow L_{E_0}E_1 \longrightarrow \text{Hom}_{\mathbf{D}}^{\bullet}(E_0, E_1) \otimes E_0 \longrightarrow E_1 \longrightarrow L_{E_0}E_1[1] \longrightarrow \dots$$

The right mutation is a pair $(E_1, R_{E_1}E_0)$, where $R_{E_1}E_0$ is defined to be a cone of the triangle

$$(2.6) \quad \dots \longrightarrow R_{E_1}E_0[-1] \longrightarrow E_0 \longrightarrow \text{Hom}_{\mathbf{D}}^{\bullet}(E_0, E_1)^* \otimes E_1 \longrightarrow R_{E_1}E_0 \longrightarrow \dots$$

More generally, if $(E_0, \dots, E_n)$ is an exceptional collection of arbitrary length in $\mathbf{D}$ then one can define left and right mutations of an object $E \in \mathbf{D}$ through the category $\langle E_0, \dots, E_n \rangle$. Denote $L_{\langle E_0, \dots, E_n \rangle} E$ and $R_{\langle E_0, \dots, E_n \rangle} E$ left and right mutations of E through $\langle E_0, \dots, E_n \rangle$, respectively. One proves ([3], Proposition 2.1) that mutations of an exceptional collection are exceptional collections.

Let $(E_0, \dots, E_n)$ be an exceptional collection in $\mathbf{D}$. One can extend it to an infinite sequence of objects (E_i) of $\mathbf{D}$, where $i \in \mathbb{Z}$, defined inductively by putting

$$(2.7) \quad E_{i+n+1} := R_{\langle E_{i+1}, \dots, E_{i+n} \rangle} E_i, \quad E_{i-n-1} := L_{\langle E_{i-n}, \dots, E_{i-1} \rangle} E_i.$$

Let X be a variety of dimension m , and ω_X the canonical invertible sheaf on X .

Definition 2.6. A sequence of objects (E_i) of $\mathbf{D}^b(X)$, where $i \in \mathbb{Z}$, is called a *helix of period n* if $E_i = E_{i+n} \otimes \omega_X[m - n + 1]$. An exceptional collection $(E_0, \dots, E_n)$ in $\mathbf{D}^b(X)$ is called a *thread of the helix* if the infinite sequence (E_i) , obtained from the collection $(E_0, \dots, E_n)$ as in (2.7), is a helix of period $n + 1$.

For a Fano variety X there is a criterion to establish whether a given exceptional collection $(E_0, \dots, E_n)$ is complete, i.e. generates $\mathbf{D}^b(X)$.

Theorem 2.1 ([3], Theorem 4.1). *Let X be a Fano variety, and $(E_0, \dots, E_n)$ be an exceptional collection in $\mathbf{D}^b(X)$. The following conditions are equivalent:*

- (1) *The collection $(E_0, \dots, E_n)$ generates $\mathbf{D}^b(X)$.*
- (2) *The collection $(E_0, \dots, E_n)$ is a thread of the helix.*

3. A THEOREM ON SEMIORTHOGONAL DECOMPOSITIONS

The theorem below is a generalization of Orlov's theorem on semiorthogonal decompositions in the derived categories of projective bundles ([17], Theorem 2.6).

Theorem 3.1. *Let $\pi : X \rightarrow S$ be a flat proper morphism between two smooth schemes X and S . Let $\mathcal{E}_1, \dots, \mathcal{E}_n$ be a set of objects of $\mathbf{D}^b(X)$ such that:*

- (i) *For any point $s \in S$ the restrictions $\mathcal{E}_i^s := \mathcal{E}_i \otimes \mathcal{O}_{X_s}$, X_s being the fiber $\pi^{-1}(s)$, satisfy semiorthogonality condition, that is for $i > j$ one has $\text{Hom}_{X_s}^\bullet(\mathcal{E}_i^s, \mathcal{E}_j^s) = 0$.*
- (ii) *For $i = 1, \dots, n$ and any point $s \in S$ the object $\mathcal{E}_i^s$ is an exceptional object of $\mathbf{D}^b(X_s)$.*
- (iii) *For any $s \in S$ the collection $(\mathcal{E}_1^s, \dots, \mathcal{E}_n^s)$ generates $\mathbf{D}^b(X_s)$.*

Put it simply, the above conditions say that for any $s \in S$ the collection of objects $\mathcal{E}_1^s, \dots, \mathcal{E}_n^s$ is a complete exceptional collection in $\mathbf{D}^b(X_s)$.

Then $\mathbf{D}^b(X)$ has a semiorthogonal decomposition:

$$(3.1) \quad \mathbf{D}^b(X) = \langle \pi^*(\mathbf{D}^b(S)) \otimes \mathcal{E}_1, \dots, \pi^*(\mathbf{D}^b(S)) \otimes \mathcal{E}_n \rangle$$

Here $\pi^(\mathbf{D}^b(S)) \otimes \mathcal{E}_i$ denotes the full triangulated subcategory in $\mathbf{D}^b(X)$ generated by objects $\pi^*(\mathcal{F}) \otimes \mathcal{E}_i$, $\mathcal{F} \in \mathbf{D}^b(S)$. Each subcategory $\pi^*(\mathbf{D}^b(S)) \otimes \mathcal{E}_i$ in the decomposition (3.1) is equivalent to $\mathbf{D}^b(S)$.*

Proof. For $i = 1, \dots, n$ define functors $\Phi_i : \mathbf{D}^b(S) \rightarrow \mathbf{D}^b(X)$ by putting $\Phi_i(A) = \pi^*(A) \otimes \mathcal{E}_i$, where $A \in \mathbf{D}^b(S)$. We check first that each Φ_i is a full faithful embedding of $\mathbf{D}^b(S)$ into $\mathbf{D}^b(X)$.

Lemma 3.1. *For $i = 1, \dots, n$ the functors $\Phi_i : \mathbf{D}^b(S) \rightarrow \mathbf{D}^b(X)$ are full faithful embeddings.*

Proof. Let $A, B \in D^b(S)$. For each $i = 1, \dots, n$ one has to show that

$$(3.2) \quad \text{Hom}_X^\bullet(\Phi_i(A), \Phi_i(B)) = \text{Hom}_S^\bullet(A, B).$$

By definition of the functors Φ_i , the left-hand side of (3.2) is isomorphic to

$$(3.3) \quad \text{Hom}_X^\bullet(\pi^*(A) \otimes \mathcal{E}_i, \pi^*(B) \otimes \mathcal{E}_i) = \text{Hom}_S^\bullet(A, B \otimes \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)).$$

The isomorphism (3.3) follows from adjunction of the functors π^* and π_* , and projection formula. Let us show that $\pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)$ is quasiisomorphic to $\mathcal{O}_S$. To this end, consider the diagram (cartesian square):

$$\begin{array}{ccc} X_s & \xrightarrow{\tilde{i}_s} & X \\ \pi_s \downarrow & & \downarrow \pi \\ S & \xrightarrow{i_s} & S \end{array}$$

Since the morphism π is flat, the base change holds ([11], Corollary 2.23):

$$(3.4) \quad \pi_{s*} \tilde{i}_s^* = i_s^* \pi_*$$

Hence,

$$(3.5) \quad \mathbb{H}^*(X_s, \tilde{i}_s^* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)) = \pi_{s*} \tilde{i}_s^* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i) = i_s^* \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i).$$

By assumption (ii) of the theorem, for any $s \in S$ one has

$$(3.6) \quad \mathbb{H}^*(X_s, \tilde{i}_s^* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)) = \mathbb{H}^*(X_s, \text{Hom}_{X_s}^\bullet(\mathcal{E}_i^s, \mathcal{E}_i^s)) = \mathbf{k}.$$

Thus, for any $s \in S$

$$(3.7) \quad i_s^* \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i) = \mathcal{O}_s.$$

Further, one has a canonical morphism $c: \mathcal{O}_X \xrightarrow{\tilde{c}} \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)$. Note that $\mathcal{O}_X = \pi^* \mathcal{O}_S$. Thus, by adjunction one arrives at a distinguished triangle:

$$(3.8) \quad \dots \longrightarrow \mathcal{O}_S \xrightarrow{\tilde{c}} \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i) \longrightarrow \text{Cone}(\tilde{c}) \longrightarrow \mathcal{O}_S[1] \longrightarrow \dots$$

Applying the functor i_s^* to the triangle (3.8) and taking into account (3.6), and (3.7), one gets

$$(3.9) \quad i_s^* \text{Cone}(\tilde{c}) = 0$$

for any $s \in S$. Indeed, by flatness of π , the morphism $i_s^*(\tilde{c}): \mathcal{O}_s \rightarrow i_s^* \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i)$, obtained from the triangle (3.8), is the isomorphism in (3.6). Recall that the set $\Omega_S = \{\mathcal{O}_s : s \in S\}$, where s ranges over the set of all points in S , is a spanning class for $D^b(S)$ ([6], Example 2.2). It follows (*loc.cit.*) that an object $\mathcal{F}$ of $D^b(S)$ such that $\text{Hom}_S^\bullet(\mathcal{F}, \mathcal{O}_s) = 0$ for all $s \in S$ is quasiisomorphic to zero. For any $\mathcal{F} \in D^b(S)$ one has an isomorphism

$$(3.10) \quad i_s^* \mathcal{F} = \text{Hom}_S^\bullet(\mathcal{F}, \mathcal{O}_s)^\vee.$$

Taking $\mathcal{F}$ to be equal to $\text{Cone}(\tilde{c})$ in (3.10) and using (3.9), one gets $\text{Cone}(\tilde{c}) = 0$. Coming back to (3.8) one sees that $\pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, \mathcal{E}_i) = \mathcal{O}_S$, q.e.d. $\square$

Put $D_i := \Phi_i(D^b(S)) \subset D^b(X)$. Let us check that each subcategory D_i is admissible in $D^b(X)$.

Lemma 3.2. *The subcategories $D_1, \dots, D_n \subset D^b(X)$ are admissible.*

Proof. One has to make sure that the functors $\Phi_i: D^b(S) \hookrightarrow D^b(X)$ have left and right adjoints. Indeed, for $A \in D^b(S)$ and $B \in D^b(X)$ one has

$$(3.11) \quad \text{Hom}_X^\bullet(\Phi_i(A), B) = \text{Hom}_X^\bullet(\pi^*(A) \otimes \mathcal{E}_i, B) = \text{Hom}_S^\bullet(A, \pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, B)),$$

where the last isomorphism in (3.11) follows from adjunction of the functors π^* and π_* , and of the functors $\otimes$ and Hom . Hence, a right adjoint functor $\Phi_i^!$ to Φ_i is isomorphic to $\pi_* \text{Hom}_X^\bullet(\mathcal{E}_i, -)$. One checks similarly that a left adjoint functor Φ_{i*} to Φ_i is isomorphic to $\pi_* \text{Hom}_X^\bullet(\mathcal{E}_i \otimes \omega_{X/S}^{-1}, -)[\dim X - \dim S]$, where $\omega_{X/S}$ is the relative canonical invertible sheaf. $\square$

We can prove now Theorem 3.1. Let us verify first the semiorthogonality condition, i.e. that for any k, l , $k > l$ one has

$$(3.12) \quad \text{Hom}_X^\bullet(D_k, D_l) = \text{Hom}_X^\bullet(\pi^*(D^b(S)) \otimes \mathcal{E}_k, \pi^*(D^b(S)) \otimes \mathcal{E}_l) = 0.$$

Indeed, by adjunction of the functors π^* and π_* this is equivalent to

$$(3.13) \quad \pi_* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l) = 0.$$

Let $s \in S$ be a point. Using once again the base change along the diagram, one gets:

$$(3.14) \quad \mathbb{H}^*(X_s, \tilde{i}_s^* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l)) = \pi_{s*} \tilde{i}_s^* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l) = i_s^* \pi_* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l).$$

By assumption (ii) of the theorem, the left-hand side of (3.14) is equal to zero. Therefore, for any $s \in S$ one has:

$$(3.15) \quad i_s^* \pi_* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l) = 0.$$

As above one gets $\pi_* \text{Hom}_X^\bullet(\mathcal{E}_k, \mathcal{E}_l) = 0$, the set Ω_S being a spanning class for $D^b(S)$. Hence, semiorthogonality is proven.

To prove that the semiorthogonal collection of subcategories $D_1, \dots, D_n$ generates $D^b(X)$ it is sufficient to show that $\langle D_1, \dots, D_n \rangle$ contains all the objects $\mathcal{O}_x$, $x \in X$, the set $\Omega_X = \{\mathcal{O}_x : x \in X\}$ being a spanning class for $D^b(X)$. Every point $x \in X$ lies in the fiber $X_s = \pi^{-1}(s)$ for some $s \in S$. By assumption (iii) of the theorem it follows that the sheaf $\mathcal{O}_x$ belongs to the category generated by the objects $\mathcal{E}_i^s$:

$$(3.16) \quad \mathcal{O}_x \in \langle \mathcal{E}_1^s, \dots, \mathcal{E}_n^s \rangle.$$

Recall that $\mathcal{E}_i^s = \mathcal{E}_i \otimes \mathcal{O}_{X_s}$ by definition. Since $\pi^*(\mathcal{O}_s) = \mathcal{O}_{X_s}$, the sheaf $\mathcal{O}_x$ belongs to $\langle D_1, \dots, D_n \rangle$ for any $x \in X$, therefore this semiorthogonal collection contains the spanning class Ω_X . This completes the proof. $\square$

An immediate corollary to Theorem 3.1 is a well-known fact:

Corollary 3.1. *Let X and Y be two smooth projective varieties. Assume that $(E_1, \dots, E_n)$ and $(F_1, \dots, F_m)$ are complete exceptional collections in $D^b(X)$ and $D^b(Y)$, respectively. Then $(E_1 \boxtimes F_1, \dots, E_n \boxtimes F_m)$ is a complete exceptional collection in $D^b(X \times Y)$.*

4. COMPLETE EXCEPTIONAL COLLECTIONS ON FLAG VARIETIES

In this section we show that the derived categories of flag varieties of classical type are generated by complete exceptional collections.

Theorem 4.1. *Let $\mathbf{G}$ be a semisimple algebraic group of classical type, $\mathbf{B}$ a Borel subgroup, and $\mathbf{G}/\mathbf{B}$ the flag variety of $\mathbf{G}$. Then $D^b(\mathbf{G}/\mathbf{B})$ is generated by an exceptional collection.*

Proof. Recall first that the case of homogeneous spaces of the groups of type $\mathbf{A}_n$ was treated by M. Kapranov ([9], Theorem 3.10). We restrict ourselves, therefore, to the groups of types $\mathbf{B}_n$, $\mathbf{C}_n$, and $\mathbf{D}_n$. The proof is based on parabolic induction (e.g., [15]), and we refer the reader to *loc.cit.* for details of this construction. Let us start with the symplectic group $\mathbf{Sp}_{2n}$, which corresponds to the type $\mathbf{C}_n$, this case being the simplest. Fix a symplectic vector space V of dimension $2n$, i.e. a space equipped with a non-degenerate skew form $\omega \in \wedge^2 V^*$. The homogeneous space, which corresponds to the leftmost node of the Dynkin diagram of $\mathbf{C}_n$, is isomorphic to $\mathbb{P}(V) = \mathbb{P}^{2n-1}$. Homogeneous spaces of the group $\mathbf{Sp}_{2n}$ are varieties of partial isotropic flags in V with respect to ω . The variety of complete isotropic flags IF_V , which is isomorphic to $\mathbf{Sp}_{2n}/\mathbf{B}$, is equipped with a set of universal bundles $\mathcal{U}_i$, where $i = 1, \dots, n$, fitting into a sequence:

$$(4.1) \quad 0 \subset \mathcal{U}_1 \subset \mathcal{U}_2 \subset \dots \subset \mathcal{U}_n = \mathcal{U}_n^\perp \subset \mathcal{U}_{n-1}^\perp \subset \dots \subset \mathcal{U}_1^\perp \subset V \otimes \mathcal{O}_{\mathrm{IF}_V}.$$

Here $\mathcal{U}_i^\perp$ is a vector bundle that is orthogonal to $\mathcal{U}_i$ with respect to ω . The flag variety IF_V can be obtained as an iteration of projective bundles over $\mathbb{P}(V)$. Indeed, for $m \leq n$ denote $\mathrm{IF}_{1,2,\dots,m}$ the partial flag variety of isotropic flags of type $(1, 2, \dots, m)$, the variety IF_V being isomorphic to $\mathrm{IF}_{1,2,\dots,n}$. By forgetting the top index, the variety $\mathrm{IF}_{1,2,\dots,m}$ becomes the projectivization of symplectic vector bundle $\mathcal{N}_{m-1} = \mathcal{U}_{m-1}^\perp / \mathcal{U}_{m-1}$ over $\mathrm{IF}_{1,2,\dots,m-1}$. Arguing by induction on m , the base of induction being $\mathbb{P}^{2n-1} = \mathrm{IF}_1$, and applying Beilinson's theorem ([1]), and Theorem 3.1, one obtains complete exceptional collections on homogeneous spaces $\mathrm{IF}_{1,2,\dots,m}$ for $m \leq n$. For convenience of the reader, let us explicitly state the answer for the flag variety IF_V . Denote $\pi_m: \mathrm{IF}_{1,2,\dots,m+1} \rightarrow \mathrm{IF}_{1,2,\dots,m}$ the projection, π_0 being the projection to the point, and p_m the composition morphism $\pi_m \circ \dots \circ \pi_{n-1}: \mathrm{IF}_{1,2,\dots,n} \rightarrow \mathrm{IF}_{1,2,\dots,m}$. Let $\mathcal{O}_{\mathcal{N}_{m-1}}(-1)$ be the Grothendieck line bundle on $\mathbb{P}_{\mathrm{IF}_{1,2,\dots,m-1}}(\mathcal{N}_{m-1}) = \mathrm{IF}_{1,2,\dots,m}$. Then the collection of line bundles

$$(4.2) \quad (\mathcal{O}_{\mathcal{N}_{n-1}}(j_{n-1}) \otimes p_{n-1}^* \mathcal{O}_{\mathcal{N}_{n-2}}(j_{n-2}) \otimes \dots \otimes p_m^* \mathcal{O}_{\mathcal{N}_{m-1}}(j_{m-1}) \otimes \dots \otimes p_1^* \mathcal{O}_{\mathbb{P}^{2n-1}}(j_0))$$

is a complete exceptional collection in $D^b(\mathrm{IF}_V)$. Here the indices j_k vary in the intervals $J_k := \{-2n + 2k < j_k \leq 0\}$, the ordering on the set of indices $(j_{n-1}, \dots, j_0)$ being the product of linear orders on J_k .

Consider now the case of orthogonal groups which correspond to the types $\mathbf{B}_n$ and $\mathbf{D}_n$. Fix an orthogonal vector space V of dimension equal to either $2n + 1$ or $2n$ for the groups of types $\mathbf{B}_n$ and $\mathbf{D}_n$, respectively. The vector space V is equipped with a non-degenerate symmetric form $q \in S^2 V^*$. The homogeneous spaces, which correspond to the leftmost nodes of the Dynkin diagrams of $\mathbf{B}_n$ and $\mathbf{D}_n$, are smooth quadrics in $\mathbb{P}(V)$ of dimensions $2n - 1$ and $2n - 2$, respectively. The derived category of a quadric was studied in [9]. Recall its description:

Theorem 4.2 ([9], Theorem 4.10). *Let $Q \subset \mathbb{P}^{n+1}$ be a smooth quadric of dimension n . Then $D^b(Q)$ is generated by an exceptional collection*

$$(4.3) \quad \Sigma(-n), \mathcal{O}_Q(-n+1), \dots, \mathcal{O}_Q(-1), \mathcal{O}_Q$$

if n is odd, and

$$(4.4) \quad \Sigma_+(-n), \Sigma_-(-n), \mathcal{O}_Q(-n+1), \dots, \mathcal{O}_Q(-1), \mathcal{O}_Q$$

if n is even.

Here Σ_-, Σ_+ are spinor bundles (they are isomorphic to each other if n is odd and the isomorphism class is denoted by Σ).

The homogeneous spaces of orthogonal group of either type are varieties of partial isotropic flags in V with respect to q . We restrict ourselves to the case of the orthogonal group of type $\mathbf{B}_n$, the case of $\mathbf{D}_n$ being similar. The variety of complete isotropic flags OF_V , which is isomorphic to $\mathbf{SO}_{2n+1}/\mathbf{B}$, is equipped with a set of universal bundles $\mathcal{W}_i$, where $i = 1, \dots, n$, fitting into a sequence:

$$(4.5) \quad 0 \subset \mathcal{W}_1 \subset \mathcal{W}_2 \subset \dots \subset \mathcal{W}_n \subset \mathcal{W}_n^\perp \subset \mathcal{W}_{n-1}^\perp \subset \dots \subset \mathcal{W}_1^\perp \subset V \otimes \mathcal{O}_{\mathrm{IF}_V}.$$

Here $\mathcal{W}_i^\perp$ is a vector bundle that is orthogonal to $\mathcal{W}_i$ with respect to q . An argument, which is analogous to that outlined for the group $\mathbf{Sp}_{2n}$, shows that the flag variety OF_V can be obtained as a successive iteration of smooth quadric fibrations over a smooth quadric $Q_{2n-1} \subset \mathbb{P}(V)$. Unfortunately, Theorem 3.1 cannot be applied directly to such fibrations, the reason being that spinor bundles are not globally defined in general. One could prove still that in our particular case of quadric fibrations spinor bundles do exist globally and then use Theorem 3.1. It makes little sense, however, since a general theorem on semiorthogonal decompositions for flat families of quadrics is now available ([13]). Recall briefly the setup of *loc.cit.* Let S be a smooth scheme (the base of a family), and let $\mathcal{E}$ be a vector bundle of rank n on S . Denote by $\mathcal{O}_{\mathcal{E}}(1)$ the Grothendieck invertible sheaf on $\mathbb{P}_S(\mathcal{E})$. Fix a line bundle $\mathcal{L}$ on S , and let $\varphi: \mathbf{S}^2\mathcal{E} \rightarrow \mathcal{L}$ be a symmetric bilinear form on $\mathcal{E}$ that makes $\mathcal{E}$ into an orthogonal vector bundle. The form φ defines a section of the line bundle $\mathcal{O}_{\mathcal{E}}(2) \otimes \pi^*\mathcal{L}$ on $\mathbb{P}_S(\mathcal{E})$. Let X be the zero locus of this section, i the embedding $X \hookrightarrow \mathbb{P}_S(\mathcal{E})$, and p the composition morphism $\pi \circ i: X \rightarrow S$. Finally, denote by $\mathcal{O}_{X/S}(1)$ the line bundle $i^*\mathcal{O}_{\mathcal{E}}(1)$. The projection p makes X into a quadric fibration over S . One checks, as in Theorem 3.1, that for $k = 1, \dots, n-2$ the collection of admissible subcategories $p^*D^b(S) \otimes \mathcal{O}_{X/S}(k) \subset D^b(X)$ is semiorthogonal. The right orthogonal in $D^b(X)$ to this collection is described by

Theorem 4.3 ([13], Theorem 4.2). *There exists a semiorthogonal decomposition in $D^b(X)$:*

$$(4.6) \quad D^b(X) = \langle D^b(S, \mathcal{B}_0), p^*D^b(S) \otimes \mathcal{O}_{X/S}(1), p^*D^b(S) \otimes \mathcal{O}_{X/S}(2), \dots, p^*D^b(S) \otimes \mathcal{O}_{X/S}(n-2) \rangle$$

where $D^b(S, \mathcal{B}_0)$ is the derived category of coherent sheaves of $\mathcal{B}_0$ -modules over S .

Here $\mathcal{B}_0$ is the sheaf of even parts of Clifford algebras on S (*loc.cit.*, Section 3.1).

To complete our proof, we argue again by induction, the base of induction being the quadric Q_{2n-1} . Indeed, for $m \leq n$ denote $\mathrm{OF}_{1,2,\dots,m}$ the partial flag variety of isotropic flags of type $(1, 2, \dots, m)$, the flag variety $\mathrm{OF}_{1,2,\dots,n}$ being isomorphic to OF_V . Denote $\pi_m: \mathrm{OF}_{1,2,\dots,m+1} \rightarrow$

$\text{OF}_{1,2,\dots,m}$ the projection, $\pi_0: \mathbb{Q}_{2n-1} \rightarrow \text{pt}$ being the projection to the point, and let p_m be the composition morphism $\pi_m \circ \dots \circ \pi_{n-1}: \text{OF}_{1,2,\dots,n} \rightarrow \text{OF}_{1,2,\dots,m}$. The variety $\text{OF}_{1,2,\dots,m}$ becomes then a smooth quadric fibration over $\text{OF}_{1,2,\dots,m-1}$. If $\mathcal{M}_{m-1} = \mathcal{W}_{m-1}^\perp / \mathcal{W}_{m-1}$ then there is an embedding $i_m: \text{OF}_{1,2,\dots,m} \subset \mathbb{P}_{\text{OF}_{1,2,\dots,m-1}}(\mathcal{M}_{m-1})$.

Theorem 4.3 furnishes a semiorthogonal decomposition of $\text{D}^b(\text{OF}_{1,2,\dots,m})$. Denote $\mathcal{B}_0^m$ the corresponding sheaf of even parts of Clifford algebras on $\text{OF}_{1,2,\dots,m-1}$. The structure of the sheaf $\mathcal{B}_0^m$ gets more simple in this case, the morphism π_{m-1} being smooth. Indeed, Proposition 3.15 of [13] ensures that the sheaf $\mathcal{B}_0^m$ is a sheaf of Azumaya algebras over $\text{OF}_{1,2,\dots,m-1}$. Moreover, $\mathcal{B}_0^m$ splits as a sheaf of Azumaya algebras, as shows the following lemma.

Lemma 4.1. *Let X be a variety admitting a cellular algebraic decomposition. Then the cohomological Brauer group $\text{Br}'(X) = \text{H}^2(X, \mathcal{O}_X^*)_{\text{tors}}$ is trivial.*

Proof. It is known that for such a variety the cycle map cl_X provides an isomorphism between Chow groups $\text{A}_*(X)$ and homology groups $\text{H}_*(X)$ ([7], Example 19.1.11). Thus, all cycles on X are algebraic and odd-dimensional integral cohomology groups of X are trivial. Considering exponential short exact sequence, one gets an exact sequence for $\text{Br}'(X)$ ([21], Proposition 1.1):

$$(4.7) \quad 0 \rightarrow A \otimes \mathbb{Q}/\mathbb{Z} \rightarrow \text{Br}'(X) \rightarrow T \rightarrow 0$$

where T is the torsion part of $\text{H}^3(X, \mathbb{Z})$, and A is the lattice of transcendental cycles, which is isomorphic to $\text{H}^2(X, \mathbb{Z})/\text{NS}(X)$, the factor group of $\text{H}^2(X, \mathbb{Z})$ by the Neron-Severi group. Given the above remarks, we get $\text{Br}'(X) = 0$. $\square$

By [8], the Brauer group $\text{Br}(X)$, which parametrizes equivalence classes of sheaves of Azumaya algebras on X , naturally injects into $\text{Br}'(X)$. Homogeneous spaces of semisimple Lie groups have algebraic cellular decompositions by virtue of the Bruhat decomposition. By Lemma 4.1, the group $\text{Br}'(\text{OF}_{1,2,\dots,m-1})$ is trivial, thus the corresponding sheaf of Azumaya algebras $\mathcal{B}_0^m$ over $\text{OF}_{1,2,\dots,m-1}$ splits, i.e. is isomorphic to $\mathcal{E}nd(\Sigma_m)$ for a vector bundle Σ_m . If $m = 1$, i.e. one has the quadric $\mathbb{Q}_{2n-1}$, then Proposition 4.4 of [9] asserts that a splitting bundle for $\mathcal{B}_0^1$ is isomorphic to the spinor bundle $\Sigma =: \Sigma_1$. In general, we have relative spinor bundles Σ_m on $\text{OF}_{1,2,\dots,m-1}$, splitting the sheaves of Azumaya algebras $\mathcal{B}_0^m$. Finally, the category $\text{D}^b(S, \mathcal{A})$ for a split Azumaya algebra $\mathcal{A}$ is equivalent to $\text{D}^b(S)$, since abelian categories $\text{Coh}(S)$ and $\text{Coh}(\mathcal{A})$ are equivalent in this case (the equivalence being given by the map $\mathcal{F} \rightarrow \text{E} \otimes \mathcal{F}$, where $\mathcal{F} \in \text{Coh}(S)$ and E is a splitting bundle for $\mathcal{A}$). Moreover, an embedding of $\text{D}^b(\text{OF}_{1,2,\dots,m-1}, \mathcal{B}_0^m) \simeq \text{D}^b(\text{OF}_{1,2,\dots,m-1})$ into $\text{D}^b(\text{OF}_{1,2,\dots,m})$ can be given by the functor $\pi_{m-1}^*(-) \otimes \Sigma_m$ ([13], Proposition 4.9). Arguing by induction on m and applying Theorems 4.2, 4.3, and the above arguments, one obtains, therefore, complete exceptional collections on homogeneous spaces $\text{OF}_{1,2,\dots,m}$, $m \leq n$. For convenience of the reader, let us explicitly state the answer for the flag variety. Let $\mathcal{O}_{\mathcal{M}_{m-1}}(-1)$ be the Grothendieck line bundle on $\mathbb{P}_{\text{OF}_{1,2,\dots,m-1}}(\mathcal{M}_{m-1})$ and let $\mathcal{O}_{\mathcal{M}_{m-1}}(-1)$ be its restriction to $\text{OF}_{1,2,\dots,m}$. Then the collection of

vector bundles

$$\begin{aligned}
 & (\Sigma_n \otimes p_{n-1}^* \Sigma_{n-1} \otimes \cdots \otimes p_m^* \Sigma_m \otimes \cdots \otimes p_1^* \Sigma_1, \dots, \\
 & p_{n-1}^* \Sigma_{n-1} \otimes \cdots \otimes p_m^* \Sigma_m \otimes p_m^* \mathcal{O}_{M_m}(j_m) \otimes \cdots \otimes p_1^* \mathcal{O}_{Q_{2n-1}}(j_0), \dots \\
 (4.8) \quad & \mathcal{O}_{M_{n-1}}(k_{n-1}) \otimes \cdots \otimes p_m^* \mathcal{O}_{M_m}(k_m) \otimes \cdots \otimes p_1^* \mathcal{O}_{Q_{2n-1}}(k_0), \dots, \mathcal{O}_{\text{OF}_V}),
 \end{aligned}$$

where $0 \leq j_m, k_m \leq 2n - 2m - 2$ and $0 \leq m \leq n - 1$, is a complete exceptional collection in $D^b(\text{OF}_V)$. Note that Σ_n is isomorphic to $\mathcal{O}_{M_{n-1}}(-1)$. The leftmost bundle in (4.8) is the tensor product of all spinor bundles Σ_i pulled back to OF_V for $i = 1, \dots, n$, and the rightmost bundle is the structural sheaf $\mathcal{O}_{\text{OF}_V}$. Terms in the middle line of (4.8) are pull-backs to $\mathcal{O}_{\text{OF}_V}$ of tensor products of the spinor bundles and line bundles. Finally, terms in the bottom line are pull-backs to $\mathcal{O}_{\text{OF}_V}$ of tensor products of line bundles. The ordering of the collection (4.8) is a natural one obtained from the construction. $\square$

Remark 4.1. Note that the collections constructed on the flag varieties for orthogonal groups do not consist entirely of line bundles. Conjecturally, there exists a complete exceptional collection of line bundles on the flag variety of a semisimple Lie group. One would like to have a uniform construction of such collections on flag varieties.

5. HOMOGENEOUS SPACES OF THE SYMPLECTIC GROUPS OF SMALL RANKS

In this section we study homogeneous spaces of Lie groups of type $\mathbf{C}_n$ for $n \leq 4$. The main result here is Theorem 5.1 where we find a complete exceptional collection in the derived category of a homogeneous space of the group $\mathbf{Sp}_6$. Theorems 5.1 and 3.1, and the main result of [19], yield complete exceptional collections in the derived categories of all homogeneous spaces of the group $\mathbf{Sp}_6$. These results also make possible to find complete exceptional collections on some homogeneous spaces of $\mathbf{Sp}_8$.

Consider the group $\mathbf{Sp}_6$. Its Dynkin diagram is of type $\mathbf{C}_3$ and there are a total of seven homogeneous spaces of $\mathbf{Sp}_6$. Fix a symplectic vector space V of dimension 6, and let ω be a non-degenerate skew form on V , $\omega \in \wedge^2 V^*$. Maximal parabolic subgroups of $\mathbf{Sp}_6$ correspond to the nodes of the diagram $\mathbf{C}_3$. The homogeneous space, corresponding to the leftmost node is $\mathbb{P}^5$, and the homogeneous space, corresponding to the rightmost node is $\text{LG}_{3,6}$, the grassmannian of lagrangian subspaces in V . The category $D^b(\text{LG}_{3,6})$ was described in [19]. Below we treat the case of homogeneous space, corresponding to the middle node of $\mathbf{C}_3$.

This is the variety $\text{IGr}_{2,6}$ of two-dimensional isotropic subspaces in V with respect to ω . It is easy to see that $\text{IGr}_{2,6}$ is a smooth hyperplane section of the grassmannian $\text{Gr}_{2,6}$. Indeed, $\text{Gr}_{2,6}$ is embedded in $\mathbb{P}(\wedge^2 V^*)$ via Plücker embedding. The form ω defines a section s_ω of the corresponding line bundle $\mathcal{O}(1)$ on $\text{Gr}_{2,6}$. The zero locus of s_ω , that is a hyperplane section of $\text{Gr}_{2,6}$ with respect to the above embedding, is the locus of two-dimensional subspaces in V such that ω vanishes on these subspaces, that is $\text{IGr}_{2,6}$. The dimension of $\text{IGr}_{2,6}$ is therefore equal to 7. Let $\text{IF}_{1,2,6}$ be

the space of pairs $(l \subset U)$, where l and U are a one-dimensional and an isotropic two-dimensional vector subspaces in V , respectively. One has a diagram (with obvious projections):

$$\begin{array}{ccc} & \text{IF}_{1,2,6} & \\ p \swarrow & & \searrow q \\ \mathbb{P}^5 & & \text{IGr}_{2,6} \end{array}$$

The projections p and q make $\text{IF}_{1,2,6}$ into a $\mathbb{P}^3$ -bundle over $\mathbb{P}^5$ and a $\mathbb{P}^1$ -bundle over $\text{IGr}_{2,6}$, respectively. More exactly, the $\mathbb{P}^3$ -bundle over $\mathbb{P}^5$ is the projectivization of the null-correlation bundle $\mathbf{N}_\omega$ of rank 4 that corresponds to the form ω ([18], §4). Recall that $\mathbf{N}_\omega$ fits into a short exact sequence of vector bundles on $\mathbb{P}^5$:

$$(5.1) \quad 0 \longrightarrow \mathbf{N}_\omega \longrightarrow \mathcal{T}_{\mathbb{P}^5}(-1) \xrightarrow{p_\omega} \mathcal{O}_{\mathbb{P}^5}(1) \longrightarrow 0,$$

where $\mathcal{T}_{\mathbb{P}^5}$ is the tangent bundle on $\mathbb{P}^5$, and the map $p_\omega \in \text{Hom}_{\mathbb{P}^5}(\mathcal{T}_{\mathbb{P}^5}(-1), \mathcal{O}_{\mathbb{P}^5}(1)) = \bigwedge^2 V^*$ corresponds to the form ω . The $\mathbb{P}^1$ -bundle over $\text{IGr}_{2,6}$ is the projectivization of the universal bundle $\mathcal{U}$ of rank 2. From the above diagram one easily finds the rank of the Grothendieck group $K^0(\text{IGr}_{2,6})$ to be equal to 12. Given such a diagram, it is natural to suppose that the direct images under the morphism q of appropriate line bundles on $\text{IF}_{1,2,6}$ may give a complete exceptional set on $\text{IGr}_{2,6}$ (cf. [19]). This method works in this case indeed.

Theorem 5.1. *$D^b(\text{IGr}_{2,6})$ is generated by the exceptional collection:*

$$(5.2) \quad \mathbf{K} = \langle \mathcal{U}(-4), \mathcal{O}_{\text{IGr}_{2,6}}(-4), S^2\mathcal{U}(-3), \mathcal{U}(-3), \mathcal{O}_{\text{IGr}_{2,6}}(-3), S^2\mathcal{U}(-2), \mathcal{U}(-2), \\ \mathcal{O}_{\text{IGr}_{2,6}}(-2), \mathcal{U}(-1), \mathcal{O}_{\text{IGr}_{2,6}}(-1), \mathcal{U}, \mathcal{O}_{\text{IGr}_{2,6}} \rangle.$$

Proof. One immediately verifies, using the Borel–Weil–Bott theorem, that the collection $\mathbf{K}$ is exceptional. We need to prove that $\mathbf{K}$ is complete, i.e. generates the derived category.

Let $\mathcal{O}_{\mathcal{U}}(-1)$ and $\mathcal{O}_{\mathbf{N}_\omega}(-1)$ be the Grothendieck invertible sheaves on $\mathbb{P}_{\text{IGr}_{2,6}}(\mathcal{U})$ and $\mathbb{P}_{\mathbb{P}^5}(\mathbf{N}_\omega)$, respectively. Denote $\omega_{\text{IF}_{1,2,6}}$ the canonical line bundle on $\text{IF}_{1,2,6}$. There are following relations between line bundles on $\text{IF}_{1,2,6}$:

- (i) $p^*\mathcal{O}_{\mathbb{P}^5}(-1) = \mathcal{O}_{\mathcal{U}}(-1)$
- (ii) $p^*\mathcal{O}_{\mathbb{P}^5}(-1) \otimes \mathcal{O}_{\mathbf{N}_\omega}(-1) = q^*\mathcal{O}_{\text{IGr}_{2,6}}(-1)$
- (iii) $\omega_{\text{IF}_{1,2,6}} = p^*\mathcal{O}_{\mathbb{P}^5}(-2) \otimes q^*\mathcal{O}_{\text{IGr}_{2,6}}(-4) = p^*\mathcal{O}_{\mathbb{P}^5}(-6) \otimes \mathcal{O}_{\mathbf{N}_\omega}(-4)$

Let $\mathcal{L}_{i,j}$ be the sheaf $p^*\mathcal{O}_{\mathbb{P}^5}(-i) \otimes \mathcal{O}_{\mathbf{N}_\omega}(-j)$. From the above relations one finds:

$$(5.3) \quad q_*(\mathcal{L}_{i,j}) = \begin{cases} S^{j-i}\mathcal{U}(-i), & j \geq i \\ 0, & j - i = -1 \\ S^{i-j-2}\mathcal{U}(-j-1)[-1], & j - i \leq -2 \end{cases}.$$

For $i = -3, \dots, 0$ denote D_i the full subcategory $p^* D^b(\mathbb{P}^5) \otimes \mathcal{O}_{N_\omega}(i) \subset D^b(\text{IF}_{1,2,6})$. There is a semiorthogonal decomposition in $D^b(\text{IF}_{1,2,6})$ with respect to the projection p :

$$(5.4) \quad D^b(\text{IF}_{1,2,6}) = \langle D_{-3}, D_{-2}, D_{-1}, D_0 \rangle.$$

To prove the theorem we need to pick up a convenient complete exceptional collection in $D^b(\text{IF}_{1,2,6})$. To this end, choose the following exceptional collections in D_i for $i = -3, \dots, 0$:

$$(5.5) \quad \begin{aligned} D_{-3} &= \langle \mathcal{O}_{\mathbb{P}^5}(-7), \mathcal{O}_{\mathbb{P}^5}(-6), \dots, \mathcal{O}_{\mathbb{P}^5}(-3), \mathcal{O}_{\mathbb{P}^5}(-2) \rangle \otimes \mathcal{O}_{N_\omega}(-3) \\ D_{-2} &= \langle \mathcal{O}_{\mathbb{P}^5}(-6), \mathcal{O}_{\mathbb{P}^5}(-5), \dots, \mathcal{O}_{\mathbb{P}^5}(-2), \mathcal{O}_{\mathbb{P}^5}(-1) \rangle \otimes \mathcal{O}_{N_\omega}(-2) \\ D_{-1} &= \langle \mathcal{O}_{\mathbb{P}^5}(-5), \mathcal{O}_{\mathbb{P}^5}(-4), \dots, \mathcal{O}_{\mathbb{P}^5}(-1), \mathcal{O}_{\mathbb{P}^5} \rangle \otimes \mathcal{O}_{N_\omega}(-1) \\ D_0 &= \langle \mathcal{O}_{\mathbb{P}^5}(-4), \mathcal{O}_{\mathbb{P}^5}(-3), \dots, \mathcal{O}_{\mathbb{P}^5}, \mathcal{O}_{\mathbb{P}^5}(1) \rangle \otimes \mathcal{O}_{N_\omega}. \end{aligned}$$

Denote L the complete exceptional collection in $D^b(\text{IF}_{1,2,6})$ thus obtained. Let M be the set of objects of $D^b(\text{IGr}_{2,6})$ obtained by applying the functor q_* to the collection L . Write this symbolically as

$$(5.6) \quad q_* L = M.$$

From (5.3) one sees that up to a shift all the sheaves of the collection K are contained in M . If the sets K and M consisted of the same elements this would prove the theorem. This is not the case, however, since two objects of M do not belong to K :

$$(5.7) \quad q_*(\mathcal{L}_{7,3}) = S^2 \mathcal{U}(-4)[-1] \quad \text{and} \quad q_*(\mathcal{L}_{-1,0}) = \mathcal{U}^*.$$

The invertible sheaves $\mathcal{L}_{7,3}$ and $\mathcal{L}_{-1,0}$ are situated at the opposite ends of the collection L . We can mutate $\mathcal{L}_{7,3}$ to the right through L . By Theorem 2.1, this mutation is given, up to a shift, by twisting the bundle $\mathcal{L}_{7,3}$ by $\omega_{\text{IF}_{1,2,6}}^{-1} = \mathcal{L}_{-6,-4}$. We obtain, therefore, the line bundle $\mathcal{L}_{1,-1}$. Such a mutation gives rise to a new thread of the helix generated by L . With respect to this thread the category $D^b(\text{IF}_{1,2,6})$ has a semiorthogonal decomposition:

$$(5.8) \quad D^b(\text{IF}_{1,2,6}) = (\langle \mathcal{L}_{-1,0}, \mathcal{L}_{1,-1} \rangle^\perp, \langle \mathcal{L}_{-1,0}, \mathcal{L}_{1,-1} \rangle).$$

Assume that the collection K is not complete, i.e. there is a non-zero left orthogonal ${}^\perp K \neq 0$. Let $\mathcal{F}$ be a non-zero object of ${}^\perp K$. Then $q^* \mathcal{F}$ belongs to the full subcategory $\langle \mathcal{L}_{-1,0}, \mathcal{L}_{1,-1} \rangle$. Indeed, for any exceptional generator $\mathcal{M}$ of the category $\langle \mathcal{L}_{-1,0}, \mathcal{L}_{1,-1} \rangle^\perp$, one has:

$$(5.9) \quad \text{Hom}_{\text{IF}_{1,2,6}}^\bullet(q^* \mathcal{F}, \mathcal{M}) = \text{Hom}_{\text{IGr}_{2,6}}^\bullet(\mathcal{F}, q_* \mathcal{M}) = 0,$$

since $q_* \mathcal{M}$ belongs to K by the very construction. Hence, there is a distinguished triangle:

$$(5.10) \quad \dots \longrightarrow V_{1,-1} \otimes \mathcal{L}_{1,-1} \longrightarrow q^* \mathcal{F} \longrightarrow V_{-1,0} \otimes \mathcal{L}_{-1,0} \longrightarrow \dots$$

Applying the functor q_* to the triangle (5.10), one gets:

$$(5.11) \quad \dots \longrightarrow V_{1,-1} \otimes \mathcal{O}_{\text{IGr}_{2,6}}[-1] \longrightarrow \mathcal{F} \longrightarrow V_{-1,0} \otimes \mathcal{U}^* \longrightarrow \dots$$

Since $\text{Hom}_{\text{IGr}_{2,6}}^\bullet(\mathcal{U}^*, \mathcal{O}_{\text{IGr}_{2,6}}) = 0$, the triangle (5.11) splits and one has an isomorphism:

$$(5.12) \quad \mathcal{F} = V_{-1,0} \otimes \mathcal{U}^* \oplus V_{1,-1} \otimes \mathcal{O}_{\text{IGr}_{2,6}}[-1].$$

The graded vector space $V_{1,-1}$ is forced to be zero, since $\mathcal{F} \in {}^\perp K$ and $\mathrm{Hom}_{\mathrm{IGr}_{2,6}}^\bullet(\mathcal{U}^*, \mathcal{O}_{\mathrm{IGr}_{2,6}}) = 0$. Therefore, $\mathcal{F}$ is isomorphic to $V_{-1,0} \otimes \mathcal{U}_2^*$. Note that $\mathrm{Hom}_{\mathrm{IGr}_{2,6}}^\bullet(\mathcal{U}^*, \mathcal{U}(-4)) = \mathbf{k}[-7]$. Taking Hom -s between $\mathcal{F}$ and various shifts of the bundle $\mathcal{U}(-4)$, and using the orthogonality condition, one sees that the graded vector space $V_{-1,0}$ is zero as well. Hence, $\mathcal{F}$ is zero. This finishes the proof. $\square$

It follows easily now that the derived category of any homogeneous space for $\mathbf{Sp}_6$ is generated by an exceptional collection. Indeed, such a homogeneous space is a variety of partial isotropic flags in V , and is obtained by giving an ordered subset S of the ordered set $(1, 2, 3)$. Denote this variety IF_S . The derived categories of IF_i for $i = 1, 2, 3$, by [1], [19], and Theorem 5.1 are generated by exceptional collections. Any other homogeneous space IF_S is mapped onto one of IF_i for $i = 1, 2, 3$, and it is easy to see that IF_S is either a $\mathbb{P}^1$ -bundle over IF_i for some i , or a $\mathbb{P}^2$ -bundle over IF_3 , or a two-step iteration of $\mathbb{P}^k$ -bundles for $k = 1$ and 2 (the last case corresponds to the flag variety for $\mathbf{Sp}_6$). Therefore, applying either Orlov's theorem ([17]) or Theorem 3.1, and the above arguments one gets complete exceptional collections in the derived categories of homogeneous spaces of the group $\mathbf{Sp}_6$.

Arguing as above, we also obtain that the derived categories of some homogeneous spaces of $\mathbf{Sp}_8$ are also generated by exceptional collections. Indeed, such a homogeneous space is a variety of partial isotropic flags in a symplectic vector space W of dimension 8, and is obtained by giving an ordered subset S of the ordered set $(1, 2, 3, 4)$ (there are 16 such varieties). Let us prove that the category $D^b(\mathrm{IF}_S)$, where the subset S contains the element 1, has an exceptional collection. Indeed, such a homogeneous space IF_S is mapped onto $\mathbb{P}(W)$. Denote IF_W the flag variety and consider first the homogeneous spaces $\mathrm{IF}_{1,3}$ and $\mathrm{IF}_{1,4}$. Let $\mathcal{U}_i$, where $i = 1, \dots, 4$, be the sequence of tautological bundles on IF_W as in Section 3:

$$(5.13) \quad 0 \subset \mathcal{U}_1 \subset \mathcal{U}_2 \subset \dots \subset \mathcal{U}_4 = \mathcal{U}_4^\perp \subset \mathcal{U}_3^\perp \subset \dots \subset \mathcal{U}_1^\perp \subset V \otimes \mathcal{O}_{\mathrm{IF}_W}.$$

Denote $\pi_{1,3}$ and $\pi_{1,4}$ the projections from $\mathrm{IF}_{1,3}$ and $\mathrm{IF}_{1,4}$ onto $\mathrm{IF}_1 = \mathbb{P}(W)$, respectively. The projections $\pi_{1,3}$ and $\pi_{1,4}$ are smooth morphisms with fibers isomorphic to $\mathrm{IGr}_{2,6}$ and $\mathrm{LG}_{3,6}$, respectively; in other words, these fibers being homogeneous spaces of $\mathbf{Sp}_6$. There is the universal symplectic bundle $\mathcal{U}_1^\perp/\mathcal{U}_1$ on both spaces $\mathrm{IF}_{1,3}$ and $\mathrm{IF}_{1,4}$. Applying Theorem 3.1, [19], and Theorem 5.1 one obtains complete exceptional collections in the categories $D^b(\mathrm{IF}_{1,3})$ and $D^b(\mathrm{IF}_{1,4})$. Any other IF_S as above is either a $\mathbb{P}^1$ -bundle, or a two-step iteration of $\mathbb{P}^1$ -bundles over either $\mathrm{IF}_{1,3}$, or $\mathrm{IF}_{1,4}$. One therefore obtains complete exceptional collections in these cases as well. Details are left to the reader.

Remark 5.1. For symplectic groups of arbitrary ranks the above arguments give that the study of derived categories of homogeneous spaces of these groups can be reduced, in a way, to that of homogeneous spaces corresponding to maximal parabolic subgroups. Indeed, if $\mathbf{P} \subset \mathbf{P}_m$ are two parabolic subgroups of $\mathbf{Sp}_{2n}$, and $\mathbf{P}_m$ is maximal parabolic, then the parabolic induction permits to map a homogeneous space $\mathbf{Sp}_{2n}/\mathbf{P}$ onto $\mathbf{Sp}_{2n}/\mathbf{P}_m$ with fibers being homogeneous spaces of either symplectic or linear groups of smaller rank. Thus, if one knows the derived categories of homogeneous spaces of groups of smaller rank and those of homogeneous spaces that correspond to maximal parabolic subgroups of $\mathbf{Sp}_{2n}$ then Theorem 3.1 furnishes semiorthogonal decompositions in the derived categories $D^b(\mathbf{Sp}_{2n}/\mathbf{P})$. The same argument is valid for homogeneous spaces of

orthogonal groups. Finally, let us remark that the construction of exceptional collections outlined in the proof of Theorem 5.1 may be extended to a larger class of homogeneous spaces. Details of the computations will be given elsewhere.

Acknowledgements. This paper owes its existence to numerous illuminating conversations with A. Kuznetsov. He also made a number of valuable remarks on a preliminary version of this text. I thank him heartily. I would like to express also my gratitude to the University Paris 13, where a part of this work was done.

REFERENCES

- [1] A. Beilinson, “Coherent sheaves on $\mathbb{P}^n$ and Problems of Linear Algebra”, *Funct. Anal. Appl.*, 12 (1978), pp. 68–69
- [2] R. Bezrukavnikov, I. Mirković and D. Rumynin, “Localization of modules for a semisimple Lie algebra in prime characteristic”, *arXiv:math.RT/0205144*
- [3] A. Bondal, “Representations of associative algebras and coherent sheaves”, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 53 (1989), no. 1, pp. 25–44; translation in *Math. USSR Izv.* 34 (1990), no. 1, pp. 23–42
- [4] A. Bondal, M. Larsen and V. Lunts, “Grothendieck ring of pretriangulated categories”, *Int. Math. Res. Not.*, (2004), no. 29, pp. 1461–1495
- [5] A. Bondal and D. Orlov, “Derived categories of coherent sheaves”, *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)* (Beijing), Higher Ed. Press, 2002, pp. 47–56
- [6] T. Bridgeland, “Equivalences of triangulated categories and Fourier–Mukai transforms”, *Bull. London Math. Soc.* v. 31 (1999), no. 1, pp. 25–34
- [7] W. Fulton, “Intersection Theory”, Springer-Verlag, Berlin, New York, 1984
- [8] A. Grothendieck, “Le groupe de Brauer”, *Dix exposés sur la cohomologie des schémas*, (1968), pp. 46–198, North-Holland, Amsterdam
- [9] M. Kapranov, “On the derived category of coherent sheaves on some homogeneous spaces”, *Inv. Math.*, 92 (1988), pp. 479–508
- [10] M. Kontsevich, “Homological algebra of mirror symmetry”, *Proceedings of the International Congress of Mathematicians, Vol. 1,2 (Zürich, 1994)* (Basel), Birkhäuser, 1995, pp. 120–139
- [11] A. Kuznetsov, “Hyperplane sections and derived categories”, (Russian) *Izv. RAN Ser. Mat.* 70 (2006), no. 3, pp. 23–128; translation in *Izv. Math.* 70 (2006), no. 3, pp. 447–547
- [12] A. Kuznetsov, “Homological projective duality”, *arXiv:math.AG/0507292*
- [13] A. Kuznetsov, “Derived categories of quadric fibrations and intersections of quadrics”, *arXiv:math.AG/0510670*
- [14] A. Kuznetsov, “Exceptional collections for Grassmannians of isotropic lines”, *arXiv:math.AG/0512013*
- [15] V. Lakshmibai, C. Musili and C.S.Seshadri, “Cohomology of line bundles on G/B ”, *Annales Scientifiques de l’É.N.S., 4^e série, tome 7, no. 1* (1974), pp. 89 – 137
- [16] S. Mukai, “Biregular classification of Fano threefolds and Fano manifolds of coindex 3”, *Proc. Nat. Acad. Sci., USA*, 86 (1989), pp. 3000–3002
- [17] D. Orlov, “Projective bundles, monoidal transformations, and derived categories of coherent sheaves”, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 56 (1992), pp. 852–862; translation in *Math. USSR Izv.* 38 (1993), pp. 133–141
- [18] C. Okonek, M. Schneider and H. Spindler, “Vector bundles on complex projective spaces”, Birkhäuser, Boston, 1980
- [19] A. Samokhin, “The derived category of coherent sheaves on LG_3^C ”, (Russian) *Uspekhi Math. Nauk*, 56, 3, (2001), pp. 592–594; translation in *Russ. Math. Surv.* 56 (2001), 3, pp. 592–594
- [20] A. Samokhin, “Tilting bundles on some Fano varieties via the Frobenius morphism”, in preparation
- [21] S. Schroer, “Topological methods for complex-analytic Brauer groups”, *Topology*, 44, (2005), pp. 875–894

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, MOSCOW, RUSSIA

E-mail address: samokhin@mccme.ru